



# **PRIMEVET**

**THE TOP CHOICE in Vet Meds!**

**Prime Veterinary Philippines, Inc.**







## A PROFILE PRIME VETERINARY PHILIPPINES, INC.

### ORGANIZATION

Prime Veterinary Philippines, Inc. (PRIMEVET) was organized in 1996 to address the growing need for top-quality and affordable veterinary products among local poultry and livestock raisers.

### PRODUCTS & SERVICES

PRIMEVET supplies hog and poultry veterinary products, such as premixes, injectables, water solubles, oral suspensions, pig doser and disinfectants to local livestock and poultry producers.

### VISION

To provide high quality and affordable veterinary medicines for every commercial swine and poultry raiser.

### MISSION

To be consistent with the credibility and reliability of all our products.



## DEXXFLOR 40% LA FLORFENICOL + DEXAMETHASONE

## LONG-ACTING ANTI-INFECTIVE "WONDER DRUG"

### FORMULATION

Each ml contains:  
Florfenicol.....400 mg  
Dexamethasone .....1 mg

### DOSAGE AND ADMINISTRATION


GROWER PIGS: 1 ml per 25 kg of body weight.  
One injection every 2 days until symptoms disappear.

By deep intramuscular (IM) injection, a dose for 48 hours interval.

### INDICATION

Indicated for the prevention and treatment of acute system diseases caused by:

- **PATHOGENS**
  - A. Pasteurella and Haemophilus in Swine and ruminants;
- **SWINE PNEUMONIC COMPLEX CAUSED BY**
  - A. Mycoplasma hypopneumoniae
  - B. Actinobacillus pleuronomoniae (APP)
  - C. MIRD Complex
  - D. Leptospirosis



Normal Lungs

### LUNGS WITH ACTINOBACILLUS PLEUROPNEUMONIAE (APP) MAY LOOK LIKE

- In acute cases,
- the lungs are **DARK AND SWOLLEN AND OOZE BLOODY FLUID** from the cut surface;
  - Hemorrhagic, even necrotic, bullae of various sizes maybe present



Source: google/yahoo





### FENDESIN

#### FORMULATION

Each ml contains:  
Tylosin (as tartrate) ..... 50 mg  
Florfenicol ..... 100 mg  
Dexamethasone ..... 1 mg

#### DOSAGE AND ADMINISTRATION

By deep intramuscular (IM) injection, once a day for 3-5 days

**SMALL ANIMALS:** 1 ml per 10 kg body weight

**LARGE ANIMALS:** 1 ml per 15-20 kg body weight

#### INDICATIONS

For the treatment of complex bacterial infectious and inflammatory disease processes sensitive to Tylosin and Florfenicol such as

- Mycoplasma-Included Respiratory Disease (MIRD) complex,
- APP Pneumonia,
- Genitourinary tract infections,
- Bacterial enteritis,
- Soft tissue infections
- Other diseases requiring combined antibiotic
- Corticosteroid therapy in swine, cattle and other livestock

### TYLOPRIME - 200

#### FORMULATION

Each ml contains:  
Tylosin base ..... 200 mg

#### DOSAGE & ADMINISTRATION

By intramuscular (IM) infection.

PIGS: 1 mL per 20 kg body weight until 24 hours after remission of the disease signs, but not to exceed 3 days

POULTRY: 0.1 to 0.2 mL per bird, subcutaneously

#### INDICATION

Intramuscular (IM) injection of tylosin can be given to pigs and cattle.

- It is used in the treatment of swine dysentery associated with:
  - a. *Vibrio coli* and/or *Treponema hyodysenteriae*
  - b. Pasteurellosis
  - c. Contagious agalactiae
  - d. Foot rot.
- It is specifically active against *Mycoplasma gallisepticum* S6 which is implicated in the Chronic Respiratory Disease (CRD) complex in chicken and infectious sinusitis in turkey.
- Among poultry, Tylosin is administered subcutaneously.



### ENROPRIME

#### FORMULATION

Each ml contains:  
Enrofloxacin ..... 100 mg

#### DOSAGE AND ADMINISTRATION

General dose rate: 2.5 mg/kg body weight

PIGS: 1 ml / 20 kg body weight; Inject IM via ham or neck muscle

Treatment to be given once a day for 3 to 5 consecutive days

#### INDICATION

For the treatment of genitor-urinary tract and respiratory infections, serious infections such as septicemia and pneumonia, skin and many soft tissue infection and intra abdominal infection

### PRIMULIN-100

#### FORMULATION

Each ml contains:  
Tiamulin (as hydrogen Fumarate)..... 100 mg

#### DOSAGE AND ADMINISTRATION

To be administered by intramuscular injection (IM)  
SWINE: 1 mL per 10 kg body weight to be given for 5 consecutive days

#### INDICATION

- For the treatment of disease/infections caused by Gram-positive organism, mycoplasma anaerobes.
- Tiamulin is used largely in swine to treat swine dysentery, enzootic pneumonia caused by:
  - a. *Mycoplasma hyopneumoniae*
  - b. *Haemophilus suis*
  - c. *Mycoplasma hyorhinis* and other pathogens
  - d. *Mycoplasma arthritis* (*Mycoplasma hyosynoviae*).



### IRON DEXTRAN



#### FORMULATION

##### DEXTRAPRIME 100

Each ml contains:  
Elemental Iron ..... 100 mg

##### DEXTRAPRIME 100 + B12

Each ml contains:  
Elemental Iron ..... 100 mg  
Cyanocobalamin ..... 100 mcg

##### DEXTRAPRIME 200

Each ml contains:  
Elemental Iron ..... 200 mg

#### INDICATION

Aids in the correction of anemia in animals as a result of hemorrhage, parasitism, infectious disease, avitaminosis, malnutrition, etc.

#### DOSAGE AND ADMINISTRATION

##### DEXTRAPRIME 100

By deep Intramuscular (IM) Injection.  
Inject 1 ml to 2 ml into the thigh muscle on the 2nd to 3rd day of life.

##### DEXTRAPRIME 100 + B12

By deep Intramuscular (IM) Injection.  
PREVENTION

PIGLET: 1 ml at 3rd and 10th day of life

GAMECOCK: 0.1 to 0.3 ml

TREATMENT

PIGLET: 2 ml

GAMECOCK: 0.2 ml to 0.5 ml

##### DEXTRAPRIME 200

By deep Intramuscular (IM) Injection.  
Inject 1 ml into the thigh muscle on the 2nd to 3rd day of life.

### CETAPRIME-B2

#### FORMULATION

Each ml contains:  
Paracetamol ..... 50mg  
Vitamin B2 ..... 10 mcg

#### INDICATION

Indicated for:

- Symptomatic relief of pain and fever in cases of bacterial infections
- Other diseases needing analgesic and antipyretic therapy.
- For the treatment and prevention of Vitamin B2 deficiency

#### DOSAGE AND ADMINISTRATION

By Intramuscular (IM) Injection, **once or twice daily for 2 – 3 days.**

SWINE : 1 ml per 10 kg body weight

Use in conjunction with anti-infective drugs is recommended.

Do not inject subcutaneously.



### PRIMECTIN 1%



#### FORMULATION

Each mL contains:  
Ivermectin ..... 10 mg

#### DOSAGE AND ADMINISTRATION

PIGS /POULTRY- 1 mL per 33 kg body weight by subcutaneous injection in the neck.

#### INDICATION

Effectively controls and treats:

- Gastrointestinal roundworms (adults & 4th stage larva)
- Lungworms (adults & 4th stage larva)
- Sucking lice
- Mites
- Cattle grubs

### CALCIOPRIME

#### FORMULATION

Each ml contains:  
Calcium Borogluconate ..... 250 mg

#### DOSAGE AND ADMINISTRATION

Pig : 25 to 50 ml (acute hypocalcemia)  
: 5 to 10 ml daily (chronic hypocalcemia)

#### INDICATION

It is indicated for the correction of calcium deficiency in cases like parturient paresis or milk fever, eclampsia, osteomalacia and other hypocalcemic conditions. It is also used in anaphylactic and allergic processes, hemorrhage and poisoning by carbon tetrachloride and chlorinated insecticides.





**SPECTILINE-BH****FORMULATION**

Each mL contains:

Spectinomycin Hydrochloride ..... 100 mg  
 Lincomycin Hydrochloride ..... 50 mg  
 Bromhexine Hydrochloride ..... 1 mg

**DOSAGE & ADMINISTRATION**

To be administered via intramuscular injection.

POULTRY: 0.2 mL/kg body weight to be given daily for 3 – 5 days

SWINE: 1 mL per 10 kg body weight for 3 to 7 days or until symptoms subside.

**INDICATION**

SPECTILINE – BH is indicated for the treatment of infections caused by organisms susceptible to lincomycin and spectinomycin such as:

- Swine enzootic pneumonia
- Bronchitis
- APP pneumonia
- Upper respiratory tract infections
- Salmonellosis
- Enteritis,
- Infectious arthritis
- Soft tissue infections

**DEXAMETH - 500****FORMULATION**

Each ml contains:

Dexamethasone.....0.5 mg

**DOSAGE AND ADMINISTRATION**

Administer Intramuscularly

PIG: 8mg to 16mg (16ml to 32ml) as a single dose

Once symptoms are controlled, doses should be reduced to minimum effect.

**INDICATION**

- Used primarily as a potent anti - inflammatory drug.
- Aids in the treatment of bovine ketosis, diagnosis of Cushing's syndrome & management of episodes of malignant hyperpyrexia
- Also possesses glucogenetic activity and can be utilized in allergic reactions, stress and non-infectious inflammatory syndrome, general and localized infections and infectious inflammatory syndrome

**PRIMETOCIN - 20****FORMULATION**

Each mL contains:

Oxytocin (synthetic) ..... 20 IU

**DOSAGE & ADMINISTRATION**

During Parturition (IM)

Sows : 1 to 2 mL

Thru IV injection: 1/2 the amount of IM doses diluted 1:10 with sterile water for injection

**INDICATION**

- During prolonged parturition where foetal position and presentation are normal, and the cervix adequately dilated.
- In cases of flaccid post-parturient uterus.
- To arrest post – parturient hemorrhage.
- As an aid in case of functional agalactia.
- As an aid in the treatment of mastitis prior to the administration of chemotherapeutics.

**BRONCHOPRIME****FORMULATION**

Each mL contains:

Bromhexine Hydrochloride ..... 3 mg

**DESCRIPTION**

Bromhexine Hydrochloride is a mucolytic agent used in the treatment of respiratory disorders associated with excessive mucus.

**DOSAGE AND ADMINISTRATION**

To be administered via intramuscular (IM) injection

**Small animals:** 1mg/kg body weight 2 times daily in 7 days.

**Swine and Cattle:** 0.1 to 0.25 mg/kg or 1ml per 10kg body weight for 3 to 7 days or until signs subside.

**INDICATION**

Bromhexine Hydrochloride is indicated for the treatment of respiratory disorders associated with severe mucus such as:

- Bronchitis
- Pneumonia
- Other respiratory syndromes in domestic animals.

**NOVA-CEFUR  
CEFTIOFUR****“FOR THE TREATMENT OF  
PNEUMONIA, FOOTROT, METRITIS”****FORMULATION**

Each 100 mL contains:

Ceftiofur Hydrochloride.....5,000 mg

**DOSAGE & ADMINISTRATION**

PIGS: 1 – 3 mL per 50 kg bodyweight, once a day for 3 days (IM)

**INDICATION**

NOVA CEFUR is a suspension of Ceftiofur Hydrochloride, a broad spectrum antibiotic indicated for the treatment of:

- Pneumonia
- Pasteurellosis
- Arthritis
- Sinusitis
- Metritis
- Footrot
- Foot infections caused by Fusobacterium necrophorum infected with bacteria, Bacteroides melaninogenicus.







## MULTIVITAMINS

### FORMULATION

Each mL contains:

Vitamin A	15,000 IU
Vitamin D3	7,500 IU
Vitamin B1	10 mg
Vitamin B2	5 mg
Vitamin B6	3 mg
Vitamin B12	20 mcg
Nicotinamide	35 mg
D-Panthenol	35 mg

### DOSAGE & ADMINISTRATION

Pigs	: 3 mL to 6 mL
Piglets	: 1 mL to 2 mL

### INDICATION

- Prevention and treatment of multiple vitamin deficiencies.
- Provides vitamins necessary for the maintenance of health and metabolism especially in cases of convalescent after bacterial and viral infections.
- Increases the resistance against stress and enhances the fertility of breeders.

## PRIMEPLEX

### FORMULATION

Each mL contains:

Thiamine Hydrochloride (Vitamin B1)	10 mg
Riboflavin (Vitamin B2)	4 mg
Pyridoxine (Vitamin B6)	4 mg
Cyanocobalamin (Vitamin B12)	10 mcg
Niacinamide (Vitamin B3)	50 mg
Panthenol (Vitamin B5)	5 mg

### DOSAGE & ADMINISTRATION

Administer intramuscularly twice a week.

SWINE	Piglets: 0.5 mL
	Adult: 3 to 5 mL

POULTRY:	0.5 mL
----------	--------


### INDICATION

- **Effective for the prevention and treatment of Vitamin B deficiencies in cases of:**

- General weakness
- Anemia
- Anorexia
- Newborn animals
- Poor growth

- **For increased requirements of B-complex vitamins**

- Growth
- Pregnancy
- Stress
- Lactation
- Convalescence



## VITAMIN ADE

### FORMULATION

Each mL contains:

Vitamin A	20,000 IU
Vitamin D3	4,500 IU
Vitamin E	5 mg

### INDICATION

It is indicated for the prevention and treatment of Vitamin A, D, and E deficiencies. Increases resistance against disease and parasitic infections. Promotes growth rate and fertility. Prevents stuntedness and infertility.

### DOSAGE & ADMINISTRATION

To be administered intramuscularly (IM) injection.

**Piglet/Dog/Cat :** 5 mL - 10 mL per head in ham or neck muscle.

**Breeder Pig/Cattle/Horse :** 15 mL - 25 mL per head in neck muscle.

**Fattener Pigs/Calves/Goat :** 15 mL - 20 mL per head in neck muscle.

Repeat dose as directed by the veterinarian.



## METAPRIME-ATP BUTAPHOSPHAN + CYANOCOBALAMIN + PYRIDOSINE + ADENOSINE TRIPHOSPHATE

**Revitalize and stimulate your animals into their PINK of health, fast!**

### FORMULATION

Each mL contains:

Butaphosphan	100 mg
Cyanocobalamin (Vit B12)	50 mcg
Pyridosine (Vit B6)	4 mg
Adenosine triphosphate	1 mg

### INDICATION

Stimulates and invigorates animals during period of stress such as:

- overcrowding
- weather changes,
- before and after farrowing
- weak neonates
- low birth-weight piglets
- underweight weanlings
- infertility and poor reproductive performance.

It increase body strength during recovery periods after infections,

- parasitism,
- malnutrition,
- inappetence,
- convulsion
- paralysis (in combination with Calcium-based preparations)
- help minimize cannibalism in pig and poultry

### PIGS



**BREEDER PIGS:**  
7 - 10 ml per head.

**GROWER PIGS:**  
5 ml per head.

**PIGLETS:**  
2 - 5 ml per head.

### COCKS



**Maintenance:**  
0.3ml (IM) Twice a month

**Conditioning:**  
0.3 ml (IM) Day 1 / Day 10 /  
3 days before the fight

To be administered by deep intramuscular or subcutaneous injection only. Inject one dose every 1 - 2 days for at least 3 consecutive injections or until signs of recovery is noticeable.





## FORMULATION

Each ml contains:  
Oxytetracycline.....200mg

## DOSAGE AND ADMINISTRATION:

Animals over 50kg: 1mL per 10kg body weight  
Animals under 50kg: 1-2mL per 10kg body weight  
Poultry: 1mL per 4-5kg body weight. By Intramuscular (IM) Injection. Once a day for 3 to 4 days.

## INDICATION

For the treatment of both systemic and local infection:

### General Organ Infections: Specific Conditions:

- Bronchopneumonia
- Bacterial enteritis
- Urinary tract infections
- Cholangitis
- Metritis
- Mastitis
- Prostatitis
- Pyodermatitis
- Infectious keratoconjunctivitis in cattle (pink eye)
- Chlamydiosis
- Heartwater, anaplasmosis, actinomycosis
- Actinobacillosis
- Nocardiosis
- Erlchiosis
- Eperythrozoonosis
- Haemobartonellosis.



## OXYPRIME 20% LA



## BENZYL PENICILLIN SODIUM

### FORMULATION

5 M IU Sterile Powder for Injection  
Each vial contains:  
Penicillin G Sodium .....5,000,000 IU

### DOSAGE AND ADMINISTRATION

To be administered via intramuscular or intravenous route.

### RECONSTITUTION

To make 10mL, add 8.7mL of water for injection

- 110,000 to 220,000 IU per 10 kg body weight treatment should be repeated at intervals of 4 to 8 hours depending on the severity of the disease.

### INDICATION

- For the treatment of diseases caused by: Gram-positive bacteria, Streptococcus pyogenes, Pneumococci, Clostridium tetani, Clostridium prefringens, Corynebacterium diphtheriae.
- Also active against Bacillus anthracis, Bacteoides, Eubacterium, Fusobacterium, Listeria monocytogenes, Staphylococci species., Actinomyces bovis, Leptospira species.

## AMOXICILLIN TRIHYDRATE + GENTAMICIN SULFATE

STERILE POWDER FOR INJECTION (IM)

### FORMULATION

Each ml contains:  
Amoxicillin (as trihydrate).....150 mg  
Gentamicin (as Sulfate).....50 mg



### INDICATION

- For the treatment of infections caused by susceptible gram-positive and gram-negative bacteria like: Pneumonia, Colibacillosis, Diarrhea, Bacterial enteritis, Mastitis, Metritis, Cutaneous abscess

### DOSAGE AND ADMINISTRATION

To be given by intramuscular injection  
General dose rate: 1 ml per 10 kg bodyweight once daily for 3 days.  
SWINE: 5 - 10 ml  
PIGLETS: 1 - 5 ml

\*For large doses, administer the dose in 2 different injection sites to favour absorption.



## GENTAMICIN SULFATE

### FORMULATION

Each mL contains:  
Gentamicin Sulfate ..... 100 mg

### DOSAGE AND ADMINISTRATION

To be administered via IM or IV injection, per 10 kg body weight of animals

Swine / Poultry: 0.5 to 3 mL

### INDICATION

- For the treatment and control of keratoconjunctivitis, septicemia and bacterial infections caused by: E. Coli, Salmonella spp., Pseudomonas aeruginosa, Rhodococcus equi, Klebsiella pneumoniae, Streptococcus spp., Alcaligenes spp., Staphylococcus spp.

## PRIMECOX 5% REFILL



### FORMULATION

Per 100 ml of Primecox 5% contains 5,000 mg Toltrazuril

### INDICATION

Primecox 5% is effective in the prevention and treatment of diarrhea in neonatal piglets caused by the coccidial parasite Isospora suis.

### FOR ORAL ADMINISTRATION PREVENTIVE:

Piglets at 3 to 5 days of age: 1ml per head  
Calves at 10 days of age: 5ml per head

### TREATMENT SINGLE DOSE

Piglets: 1ml per head  
Calves: 5ml per head

\*\*Each pump stroke is equivalent to 0.5ml. Shake well before use.

### ATTRIBUTES


- A. PRIMECOX 5% is a triazone drug.
- B. PRIMECOX 5% has a broad spectrum anticoccidial and antiprotozoal activities.
- C. PRIMECOX 5% is active against all intracellular

D. PRIMECOX 5% inhibits the nuclear division of schizonts and microgamonts and the wall-forming bodies of macrogamonts.

E. PRIMECOX 5% reduces clinical signs and oocyst excretion by a single treatment

F. PRIMECOX 5% promotes resistance to re-infection.

Source: google/yahoo



### DID YOU KNOW?

There are three types of **COCCIDIA** for pigs; **EIMERIA**, **ISOSPORA** and **CRYPTOSPORIDIA**.

And Isospora suis is the most pathogenic of the types of Coccidia.

## PRIMESCOUR



### FORMULATION

Each ml contains:  
Amoxicillin (as trihydrate) ..... 1 mg  
Neomycin (as sulphate) ..... 300 mcg  
Colistin (as sulphate) ..... 25IU  
Loperamide ..... 200 mcg  
Potassium Chloride ..... 275 mcg  
Sodium Chloride ..... 250 mcg  
Magnesium Sulfate ..... 75 mcg

### INDICATION

For the prevention and treatment of bacterial diarrhea such as colibacillosis and salmonellosis in piglets, dogs, and cat.

### DOSAGE AND ADMINISTRATION

Administer orally. 1 pump stroke is equivalent to 1 ml dispensed.

Piglets 2 weeks: 1 - 2 ml for 3 to 5 consecutive days.


Piglets 3 weeks: 2 - 3 ml for 3 to 5 consecutive days.

In severe cases, administer twice a day for the first two days of treatment.

### WITHDRAWAL PERIOD

14 days





## AMOXICILLIN 10%

### FORMULATION

Each kilogram contains:  
Amoxicillin (as trihydrate) .....100 g

### DOSAGE AND ADMINISTRATION

CATTLE / SWINE / POULTRY  
Prevention: Mix 1 kg to 1.5 kg per ton of feeds for 3 to 5 days

Treatment: Mix 2 kg to 4 kg per ton of feeds for 5 to 7 days

### INDICATION

AMOXICILLIN 10% Medicated Feed premix is a broad-spectrum antibiotic indicated for the prevention and treatment of respiratory, gastro-intestinal and genito-urinary tract infections caused by Gram-positive and Gram-negative bacteria.



## COLEXIN

### FORMULATION

Each Kg contains:  
Cefalexin monohydrate ..... 50g  
Colistin sulfate. .... 12,000,000 IU

### DOSAGE AND ADMINISTRATION

Thoroughly mix with animal feeds

### POULTRY

Prevention: Mix 500gms Colexin per ton

### INDICATION

For the prevention and treatment of moderate to severe bacterial infections involving the respiratory, gastro-intestinal, reproductive, urinary tract, as well as skin and soft tissues caused by susceptible Gram-positive and Gram-negative organisms including Mycoplasma spp.

of feeds for 3-5 days.

Treatment: Mix 1 kg Colexin per ton of feeds for 5-7 days.

### SWINE

Prevention: Mix 1 kg Colexin per ton of feeds for 3-5 days.

Treatment: Mix 1.5 – 2 kg Colexin per ton of feeds for 5-7 days.

Meat: 7 days prior to slaughter



## PRIMTIN 0.6%

### FORMULATION

Ivermectin 0.6% (medicated Powder Feed Premix)

Each kilogram contains:  
Ivermectin ..... 6 g

### INDICATION

Actively treats and controls external parasites (Sarcoptes causing Mange, Mites and Lice) as well as internal parasites (Gastrointestinal roundworms, Nodular worms, threadworms and lungworms).

### DOSAGE AND ADMINISTRATION

Growing pigs up 40 kg: 333 g/ton of feed for 7 days


40- 100 kg: 400 g/ton of feed for 7 days

Adult pigs:

- Basic limit feeding (Sows and Boars): 1.66 kg/ton of feed for 7 days

Sows on varying feed intakes:

- 200 kg average body weight (ABW)
- 2 Daily feed intakes (DFI): 1.67 kg/ton of feed for 7 days
- 2.5 DFI: 1.33 kg/ton of feed for 7 days
- 3 DFI: 1.11 kg/ton of feed for 7 days
- 250 kgs ABW 2.5 DFI: 1.67 kg/ton of feed for 7 days
- 3 DFI: 1.39 kg/ton of feed for 7 days



## PRIFENICOL 4%

Antibacterial/Anti-infective

# SAY GOODBYE TO SWINE RESPIRATORY DISEASES

- HIGH **BACTERICIDAL ACTIVITY**
- **HIGHLY EFFECTIVE** AGAINST A WIDE RANGE OF BACTERIAL INFECTIONS
- **PREVENTS AND TREATS** SWINE RESPIRATORY DISEASES

### FORMULATION

Each kilogram contains:  
Florfenicol ..... 40 g

### DOSAGE AND ADMINISTRATION

**Swine / Poultry :** 500 g – 1,000 g / ton of feed for 3-5 days or as recommended by a qualified veterinarian.

### INDICATION

- For the prevention and treatment of swine respiratory disease caused by:
  - Actinobacillus plueropneumonia
  - Pasteurella multocida
  - Streptococcus suis, Bordetella bronchiseptica
  - Haemophilus parasuis
  - Mycoplasma hyopneumoniae
- Treatment against Chronic Respiratory Disease (CRD), Coryza and Salmonellosis in poultry.



**LINCOMYCIN 4.4****FORMULATION**

Each kilogram contains:  
Lincomycin (as hydrochloride) . . . 44g

**DOSAGE AND ADMINISTRATION**

Prevention: 1 kg medicated feed premix per ton of complete feeds for 3-5 days

Treatment:

Swine dysentery: 2 kg medicate feed premix per ton of complete feeds for 5-7 days

Mycoplasmal infection: 5 kg of feed per ton of complete feed for 5-7 days

Poultry (Necrotic Enteritis): 2 kg feed per ton of complete feed for 5-7 days

**INDICATION**

For the prevention and treatment of swine disease like Dysentery and Mycoplasmal infections. In poultry (broiler chicken), It is indicated against Necrotic enteritis.

**NEOMYCIN 10%****FORMULATION**

Each kilogram contains:  
Neomycin (as Sulfate) . . . . .100 g

**INDICATION**


- Used for the prevention and treatment of bacterial enteritis:
  - a. Diarrhea
  - b. Blood dysentery
  - c. Scours
  - d. Colibacillosis
  - e. Salmonellosis
  - f. Vibronic dysentery
  - g. Diarrheas caused by Vibrio, E. coli, and Salmonella
  - h. Enterotoxemia in lambs

- Effective in the prevention and treatment of:
  - a. Salmonellosis
  - b. Bluecomb
  - c. Mud fever
  - d. Non-specific enteritis in chicken, ducks, and turkeys.

**DOSAGE AND ADMINISTRATION**

SWINE AND POULTRY

Treatment: Mix 700 g to 1.4 kg per ton of complete feed and laying hens, and 5 days for broilers.

**PARACETAMOL****FORMULATION**

Each kilogram contains:  
Paracetamol . . . . . 100,000 mg

**INDICATION**

It is indicated for symptomatic relief of pain and fever in cases of bacterial infections and other diseases needing and analgesic and antipyretic therapy.

**DOSAGE AND ADMINISTRATION**

Mix 2 kg per ton of feeds given for 3 to 4 days.

In case of severe infection, double the usual dose to increase antipyretic effect.



## TIAMULIN

Antibacterial

# FOR THE TREATMENT OF PNEUMONIA AND DYSENTERY

**FORMULATION**

Each kilogram contains:

Tiamulin Hydrogen Fumarate . . . . . 100g

**DOSAGE AND ADMINISTRATION SWINE**

Swine dysentery & enzootic pneumonia

Treatment : 1 - 1.5 kg / ton for 5-7 days.

Swine dysentery, Ileitis, Colitis:

Treatment: 1 - 1.5 kg per ton of feeds.

**POULTRY**

Treatment: 1 kg per ton of feeds

**INDICATION****SWINE**

For oral treatment of:

- Swine dysentery
- Ileitis
- Colitis
- Swine enzootic pneumonia (SEP)

**POULTRY**

For the prevention and treatment of:

- Mycoplasmal infections
- CRD (air sacculitis)
- CRD Complex
- Infectious synovitis
- Infectious coryza,
- Other bacterial sensitive to Tiamulin





## AMOX + COLISTIN WS

### FORMULATION

Each kilogram contains:  
Amoxicillin trihydrate..... 150g  
Colistin sulphate.....500,000,000 IU

### DOSAGE & ADMINISTRATION

#### SWINE/POULTRY:

- Prevention: Mix 5g (1 tsp) per 30 liters (7.5 gal) of drinking water for 3-5 days.
- Treatment: Mix 5g (1 tsp) per 15 liters (3.75 gal) of drinking water for 3-5 days.

### INDICATION

- For the treatment of gastrointestinal infections:
  - a. Colibacillosis
  - b. Salmonellosis
  - c. Bacterial enteritis
- For the treatment of respiratory infections
  - a. Pneumonia
  - b. Bronchitis
  - c. Bronchopneumonia of primary and secondary origin.
- It is also used in other bacterial infections like Mastitis, Polyarthritis and Umbilical infections in swine and infections in poultry.

## BROMITROL 10%

### FORMULATION

#### BROMITROL 10% WS

Each kilogram contains:  
Norfloxacin.....100g

### INDICATION

BROMITROL is indicated for treating gastrointestinal, respiratory and urinary tract infections caused by:

Gram-negative organisms like:

- E. coli
- Salmonella
- Haemophilus
- Pasteurella
- Treponema
- Leptospira, Bordetella
- Pseudomonas
- Campylobacter
- Vibrios

### DOSAGE & ADMINISTRATION

#### Bromitrol 10% Water Soluble Powder

#### SWINE

Prevention : 4 tsp to 20 liters of drinking water  
Treatment : 8 tsp to 20 liters of drinking water for 5 - 7 consecutive days  
Severe infections : 12 tsp to 20 liters of drinking water for 5 - 7 consecutive days.

#### POULTRY

Prevention : 2 tsp to 20 liters of drinking water  
Treatment : 4 tsp to 20 liters of drinking water for 5 - 7 consecutive days  
Severe infections: 6 tsp to 20 liters of drinking water for 5-7 consecutive days.



## CETAPRIME-100

### FORMULATION

Each kilogram contains:  
Paracetamol ..... 100g

### INDICATION

It is indicated for symptomatic relief of pain and fever in cases of bacterial infections and other diseases needing analgesic and antipyretic therapy.

### DOSAGE & ADMINISTRATION

General dose for pigs, cattle, goats, sheep, poultry:

- Dissolve 1-2 grams per liter of drinking water for 3-5 days or mix 2-4 grams per kg of feeds until recovery and fever subsides. Administer twice daily.

### RECOMMENDATION

- Prepare fresh solution daily.
- Use in conjunction with antibiotic for better treatment in cases of bacterial infections.
- Do not use in dogs and cats.
- Avoid prolonged use to prevent possible renal toxicity.
- Do not exceed recommended dose.



# MULTIVITAMINS / ELECTROLYTES



### GERLITE PLUS

- Replaces lost electrolytes in animals such as poultry, swine, calves, and foals during dehydration due to diarrhea, vomiting or during period of reduced feed intake.
- For the prevention and treatment of vitamin deficiencies.
- For Increase resistance from stress and faster recovery from diseases.

### DOSAGE AND ADMINISTRATION POULTRY

Preventive: Dissolve 5 g (1 tsp) to 10 liters (2.5gal) of drinking water continuously.

Treatment: Dissolve 5 g (1 tsp) to 4 liters (1 gal) of drinking water for 2-3 days.

### SWINE

Preventive Dissolve 5 g (1 tsp) to 4 liters (1 gal) of drinking water



### ELECTROLYTES

Provides essential electrolytes during periods of:

- Dehydration
- Stress
- Scouring vaccination
- Inadequate feed intake
- Diarrhea
- Metabolic
- Acidosis
- Ketosis

### DOSAGE AND ADMINISTRATION

To be incorporated in the drinking water.  
Add 1 tsp (5 g) to 1 gallon (4 liters) of water

### FORMULATION Each kilogram contains

VITAMINS AND MINERALS	GERLITE PLUS	ELECTROLYTES
Vitamin A	3,000,000 IU	
Vitamin D3	500,000 IU	
Vitamin E	2,000 IU	
Vitamin B1	1,000 mg	
Vitamin B2	3,000 mg	
Vitamin B5	1,000 mg	
Vitamin B6	1,000 mg	
Vitamin B12	10,000 mcg	
Vitamin K3	1,000 mg	
Folic Acid	100 mg	
Biotin	5,000 mcg	
Niacin	15,000 mg	
Niacinamide	10,000 mg	
Sodium Bicarbonate	250,000 mg	500,000 mg
Sodium Chloride	10,000 mg	255,000 mg
Potassium Chloride	20,000 mg	50,000 mg
Dextrose		195,000 mg





## BACTILLIN

### FORMULATION

Each mL contains:

Povidone Iodine . . . . . 100 mg

### INDICATIONS:

- Use for minor wounds, cuts, burns, abrasions, lacerations, prevention and disinfections of common skin infections, pre and post operative skin infection
- Infection control during insertion and care of urinary catheters, suture removal, dressing changes in circumcision, degerming during use of

intravenous devices in blood transfusion, CVP catheterization, cutdowns, disinfection of skin prior to injection or aspiration

- Infectious vaginalis, monilia, trichomonos and candida
- Hypo allergenic, non-sensitizing, non-staining
- Does not cause stinging when applied to open wounds

## PRIMECIDE

### FORMULATION

Each mL contains:

Glutaraldehyde.....150 mg

Benzalkonium Chloride.....100 mg

### INDICATION

Broad spectrum disinfectant effective against viruses, bacteria, fungi, yeast, and algae.

### APPLICATION

For general disinfection of farm facilities including animal houses, hatcheries, warehouses, transport vehicles, tire / foot dips / sprays / baths, as well as non-food surfaces in processing areas and equipment. It can also be used for aerial disinfection to reduce airborne pathogens, as water sanitizers and in flushing water lines.

Repeat disinfection every 14 days or as needed based on the degree of contamination in the affected areas.

### DILUTION & APPLICATION

**Routine Disinfection :** Dilute 10 mL in 14 liters of water.

**Water Sanitizer :** Dilute 1 mL in 2 liters of water and give continuously.

**Foot Dips / Wheel :** Dilute 10 mL in 1 liter of water

**Fogging / Aerial Disinfection :** Dilute 10 mL in 4 liters of water.

\* Repeat disinfection every 14 days or as needed based on the degree of contamination in the affected areas.



## ELIMINATOR PLUS

### All Natural Plant Fiber Extract

**ELIMINATOR PLUS (EP)** is a plant extract that is very effective in controlling disease causing agents and other harmful microorganisms such as E-coli, Salmonella, Cholera, Staphylococcus, molds, yeasts and fungi. It eliminates OFFENSIVE ODOR producing substances such as ammonia and hydrogen sulphide among others. It removes offensive smelling gases rather than mask the foul odors.

### INDICATION

- Organic, bio-degradable and 100% environment friendly
- Anti-fungal and anti-bacterial
- Instant killing action against microorganism
- Neutralizes Ammonia, Hydrogen Sulfide and other offensive odor-producing substances
- Non-toxic in its diluted form

### APPLICATIONS RECOMMENDED AND USAGE:



Concentrated (depends on count of mortality)  
Fast Decomposing Mortality, Disinfectant & Odor Eliminator



1:50 Decomposing WET Manure  
1:100 Decomposing DRY Manure



1:100 For tire bath and vehicle sprayer as disinfectant



1:100 Buildings & Facilities  
1st Application for Disinfecting & Eliminating Odor



1:200 Buildings & Facilities  
Maintenance for Disinfecting & Eliminating Odor



1:1000 Water quality enhancer for feeds and water



# CALPHOPRIME

Oral Solution Supplement



**CALPHO-PRIME OS** indicated for use in poultry for the prevention and treatment of conditions associated with mineral deficiencies, **rickets in growing birds, cage layer fatigue ,and soft shell eggs.**

**FORMULATION**  
Each kilogram contains:

Calcium Borogluconate.....	105,000mg
Phosphorus.....	100,000mg
Vitamin D3.....	180,000 IU
Sodium.....	25,000mg
Magnesium.....	6,500mg
Zinc.....	2,000mg
Copper.....	620mg
Cobalt.....	24mg
Iodine.....	10mg
Potassium.....	10mg

**BENEFITS:**

- Improve egg shell strength and hatchability
- Growth performance and feathering

**DOSAGE AND ADMINISTRATION:**  
To be given as the only source of drinking water during treatment period.

**POULTRY**

- PREVENTION: 2 ml of CALPHO-PRIME OS per liter of water to be given for two weeks.
- TREATMENT: 3 ml of CALPHO-PRIME OS per liter of water to be given immediately when signs of deficiency is observed.

## AVOID THE EFFECTS OF CALCIUM AND MINERAL DEFICIENCY:



Rickets in growing birds




Cage layer fatigue



Soft eggshells

# HEPATOGUARD

Oral Solution Supplement



STOP **FATTY LIVER SYNDROME!**

**FORMULATION**  
Each kilogram contains:

Sorbitol .....	400g
Methionine .....	25g
Vitamin B12.....	100g
Betaine.....	100g

**DOSAGE AND ADMINISTRATION**  
To be administrated via clean drinking water

**Poultry:** 1 – 2 ml per liter of drinking water, 2 - 3 consecutive days for every week for at least 3 consecutive weeks.

**Pigs:** 2 ml per liter of drinking water, 3 - 4 consecutive days for every week for at least 3 consecutive weeks.

**Breeding animals:** 2 – 3 ml per liter of drinking water, use continuously for 5 – 7 days for at least two consecutive week.

**INDICATION**

- Prevention and treatment of fatty liver syndrome, biliary, renal, and intestinal functions in poultry and other species.
- Increase the synthesis of proteins.
- Stimulates increase fertility.
- Stimulate the metabolism of sugar/carbohydrates.
- Promotion of growth and increases quality of carcass.
- For increased detoxification of the liver in cases of Mycotoxicosis and similar toxicity conditions in domesticated animals.



HEALTHY LIVER

Fatty liver in poultry can result in the following symptoms:

- Pale comb
- Performance failure
- Sunken eyes
- Mortality
- Drop in egg production
- Increased amount of abdominal fat
- Increase in body weight



FATTY LIVER

Source: google/yahoo





# PSK

## Antimalarial

### FORMULATION

Each kilogram contains:

Pyrimethamine.....	10g
Sulfamonomethoxine.....	13.2g
Vitamin K3.....	3.5g
Vitamin B12.....	0.55g


### INDICATION

Effective for the prevention and treatment of the following:

- Taxoplasmosis
- Avian Malaria
- Leukocytozoonosis
- Avian Infections
- Coryza
- Bronchitis
- Gastroenteritis
- Coccidiosis

### DOSAGE AND ADMINISTRATION

Prevention: 1 teaspoonful per 20 liters of water for 3 to 5 days  
Treatment: 1 to 1.5 tablespoons per 20 litres of water for 5 to 7 days.



# BROMITROL



### FORMULATION

#### BROMITROL 20% OS

Each kilogram contains:

Norfloxacin.....	200g
------------------	------

### INDICATION

BROMITROL is indicated for treating gastrointestinal, respiratory and urinary tract infections caused by:

Gram-negative organisms like:

- E. coli
- Salmonella
- Haemophilus
- Pasteurella
- Treponema
- Leptospira, Bordetella
- Pseudomonas
- Campylobacter
- Vibrios

### Bromitrol 20% Oral Solution

General dose is 12 mg per kg body weight.

To get a higher initial intake, withhold drinking water for 1 to 2 hours before the administration of medicated water

### Swine

Prevention: 10 ml per 20 liters of drinking water

Treatment: 20 ml per 20 liters of drinking water for 5 to 7 consecutive days

### Poultry

Prevention: 5 ml per 20 liters of drinking water

Treatment: 10ml per 20 liters of drinking water for 5 to 7 consecutive days.

# MS DRYCARE



## For optimal housing hygiene

- Active Ingredient: 90% Calcium Carbonate
- Very high absorption, >200%
- Create dry living environment so that microorganism have no chance to develop
- Easy to spread, it covers the area well and does not create dust.
- An attractive aroma - plays a role in a pleasant living environment and it can smother or neutralise unpleasant smell
- pH neutral - a hygiene powder , does not cause irritation to an animals teats, skin, hooves or legs

### PIGS



- **Piglet/sow section:**  
50 g/m<sup>2</sup>, 3 times per week
- **Farrowing pens:**  
daily, as required

### POULTRY



- **Broilers:**  
100 gr/m<sup>2</sup>, at the start of the round, and then scatter on wet patches
- **Laying nests:**  
50 g/m<sup>2</sup>, 1 time every 2 weeks





# PRIMEVET

**THE TOP CHOICE in Vet Meds!**

**Prime Veterinary Philippines, Inc.**

G/F The Esquire Center, 412 Gomezville St.  
Brgy. Addition Hills Mandaluyong City  
1552 Metro Manila Philippines

Email Add: [primevet@infarmco.com](mailto:primevet@infarmco.com)  
Cellphone No. 0998-960-6891 / 0917-509-5820  
Telephone No. (02) 8722-0413



[www.infarmco.com](http://www.infarmco.com)